



TITLE:

Vol.22表紙・総目次・Record of Works for 1970

AUTHOR(S):

CITATION:

Vol.22表紙・総目次・Record of Works for 1970. PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY 1976, 22(6)

ISSUE DATE:

1976-02-28

URL:

<http://hdl.handle.net/2433/175907>

RIGHT:

KYOTO UNIVERSITY

PUBLICATIONS
OF THE SETO MARINE
BIOLOGICAL LABORATORY

Volume XXII



PUBLISHED BY THE LABORATORY
SIRAHAMA, WAKAYAMA-KEN, JAPAN
1975-1976

CONTENTS

Nos. 1/4 (July 31, 1975)

| | |
|---|-----|
| 1. Schornikow, E. I.: Ostracod fauna of the intertidal zone in the vicinity of the Seto Marine Biological Laboratory. | 1 |
| 2. Hoestlandt, H.: Occurrences of the Isopoda Flabellifera <i>Gnorimosphaeroma rayi</i> Hoestlandt on the coasts of Japan, Eastern Siberia and Hawaii, with a brief note on its genetic polychromatism. | 31 |
| 3. Kussakin, O. G.: A list of the macrofauna in the intertidal zone of the Kurile Islands, with remarks on zoogeographical structure of the region. | 47 |
| 4. Yasuda, T.: Ecological studies on the jelly-fish, <i>Aurelia aurita</i> (Linné), in Urazoko Bay, Fukui Prefecture—XI. An observation on ephyra formation. | 75 |
| 5. Murano, M.: Mysidacea from the central and western Pacific II. Genera <i>Hyperamblyops</i> , <i>Teraterythrops</i> and <i>Synerythrops</i> (Tribe Erythropini). | 81 |
| 6. Rosell, N. C.: Two new species of barnacles (Cirripedia Thoracica) from Mindanao, Philippines. | 105 |
| 7. Baba, K. and Abe, T.: Comments on further specimens of <i>Favorinus tsuruganus</i> B. & A. from Echizen-Cho near Tsuruga Bay, Japan (Nudibranchia: Eolidoidea: Favorinidae). | 117 |
| 8. Cordero, P. A., Jr.: Some epiphytic algae in the vicinity of the Seto Marine Biological Laboratory, Wakayama Prefecture, Japan I. | 121 |
| 9. Izawa, K.: On the development of parasitic Copepoda II. <i>Colobomatus pupa</i> Izawa (Cyclopoida: Philichthyidae). | 147 |
| 10. Ross, D. M.: The behavior of pagurids in symbiotic associations with actinians in Japan. | 157 |
| 11. Morino, H.: Studies on the Talitridae (Amphipoda, Crustacea) in Japan II. Taxonomy of sea-shore <i>Orchestia</i> , with notes on the habitats of Japanese sea-shore talitrids. | 171 |
| 12. Imafuku, M.: Peristalsis in the monopolypid stage of the sea-pen, <i>Cavernularia obesa</i> Valenciennes. | 195 |
| 13. Nishikawa, T. and Tokioka, T.: Contributions to the Japanese ascidian fauna XXVI. Notes on simple ascidians commonly found in the waters near the Sabiura Marine Park Research Laboratory. | 217 |

No. 5 (November 29, 1975)

| | |
|--|-----|
| 14. Cordero, P. A., Jr.: Notes on some little-known red algal epiphytes of Japan. | 223 |
| 15. Gamô, S.: <i>Platysympus japonicus</i> sp. nov., a new cumacean Crustacea from Sagami Bay. | 229 |
| 16. Utinomi, H.: Octocorallia collected by trawling in the Western Australia (A biological result of I.I.O.E.—Collections 1963). | 237 |

| | | |
|-----|--|-----|
| 17. | Cordero, P. A., Jr.: On the occurrence of Genus <i>Actinotrichia</i> in the Philippines. | 267 |
| 18. | Baba, K.: <i>Neomenia yamamotoi</i> spec. nov., a gigantic solenogaster (Mollusca: Class Solenogastres, Family Neomeniidae), occurring in the north-eastern part of Japan. | 277 |
| 19. | Hayashi, K.: <i>Hippolysmata grabhami</i> Gordon, a synonym of <i>Lysmata amboinensis</i> (de Man) (Decapoda, Caridea, Hippolytidae). | 285 |
| 20. | Fujino, T.: Occurrence of <i>Rhynchocinetes rigens</i> Gordon, 1936 (Crustacea, Decapoda, Rhynchocinetidae) in the Indo-Pacific region. | 297 |
| 21. | Ho, J.-S.: Cyclopoid copepods of the family Chondracanthidae parasitic on New Zealand marine fishes. | 303 |
| 22. | Taniguti, T.: Droplets from the plankton net XXVI. Notes on the excretion in <i>Eurhamphaea vexilligera</i> Gegenbaur (Ctenophora). | 321 |
| 23. | Tokioka, T. and Nishikawa, T.: Contributions to the Japanese ascidian fauna XXVII. Some ascidians from Okinawa, with notes on a small collection from Hong Kong. | 323 |
| 24. | Yoshino, T.: <i>Parapercis gushikeni</i> , a new mugiloid fish from the Ryukyu Islands. | 343 |
| 25. | Uchida, T. and Sugiura, Y.: On the formation of medusa buds in <i>Proboscoidactyla ornata</i> | 347 |

No. 6 (February 28, 1976)

| | | |
|-----|---|-----|
| 26. | Furuhashi, K.: Diel vertical migration suspected in some copepods and chaetognaths in the inlet waters, with a special reference to behavioural differences between male and female, noted in the former. | 355 |
| 27. | Hamatani, I.: Preliminary account of a new species of <i>Volvatella</i> Pease, 1860, <i>V. viridis</i> sp. nov., found in the caulerpan microfauna in Japan (Opisthobranchia: Sacoglossa). | 371 |
| 28. | Nishikawa, T. and Tokioka, T.: Contributions to the Japanese ascidian fauna XXIII. Ascidians from the Amami Islands. | 377 |

Composition of the fixed sea urchin colony on Hatakezima Island, 1975.403

Bioassay data for marine pollution using sea urchin eggs, 1974.405

Oceanographic data at the S.M.B.L., 1972.409

Record of works done at the Seto Marine Biological Laboratory (published in 1970).417

RECORD OF WORKS DONE AT THE SETO MARINE BIOLOGICAL LABORATORY

1970 (昭和45年)

Contribution
No. (Jap. No.)

- *荒賀忠一：ナポリ臨海実験所における研究材料生物の供給システムについて。学術月報, **23** (1), 29—35.
[Araga, Chuichi: On the system of supplying living materials for biological researches at Stazione Zoologica di Napoli. Jap. Scientific Monthly, **23** (1), 29—35.]
- *馬場菊太郎：ムラサキクルマナマコの骨片。採と飼, **23** (1), 22.
[Baba, Kikutaro: Spicules of *Polycheira rufescens* (Brandt). Collect. & Breed., **32** (1), 22.]
- *———：日本産ヤマトウミコチョウ科およびルンキナ科種名目録。同上誌, **32** (2), 46—48.
[———： List of the Gastropteridae and Runcinidae from Japan. Ibid., **32** (2), 46—48.]
- *———：日本産アメフラシ科の種名目録。同上誌, **32** (3), 94—96.
[———： List of the Aplysiidae from Japan. Ibid., **32** (3), 94—96.]
- *———：日本産クロスジアメフラシ類追記。同上誌, **32** (11), 376—377.
[———： Additional notes on *Stylocheilus* spp. from Japan. Ibid., **32** (11), 376—377.]
- *———：日本産カノコキセワタ科の種名目録。同上誌, **32** (12), 414—415.
[———： List of the Doridiidae from Japan. Ibid., **32** (12), 414—415.]
- Baba, Kikutaro & Hamatani, Iwao: Occurrences of specimens presumably identifiable with *Stiliger ornatus* Ehrenberg, 1831, at Seto, Kii, middle Japan (Opisthobranchia: Sacoglossa). Publ. Seto Mar. Biol. Lab., **18** (3), 199—206, pls. III—IV.525
- & ———： The anatomy of *Ercolania boodleae* (Baba, 1938) from Seto, Kii, middle Japan (Opisthobranchia: Sacoglossa). Ibid., **18** (4), 215—222, pls. V—VII.528
- Bieri, Robert: Droplets from the plankton net XXIV. The predominant type of *Velella* in Japanese waters. Ibid., **17** (5), 303—304.515
- ： The food of *Porpita* and niche separation in three neuston coelenterates. Ibid. **17** (5), 305—307.516
- *布施慎一郎：砂泥底エピベントスの採集法について。
[Fuse, Shin-ichiro: On the sampling of epibenthos on sandy-muddy bottoms. Benthos Research, no. 1, 48—50.]
- Humes, A. G.: *Paramacrochiron japonicus* n. sp., a cyclopoid copepod associated with a medusa in Japan. Publ. Seto Mar. Biol. Lab., **18** (4), 223—232.529
- Illg, Paul L.: Report on ascidicole Copepoda collected during the Melanesia Expedition of the Osaka Museum of Natural History, Osaka, Japan. Ibid., **18** (3), 169—188.527

- *小林直正：ウニ受精卵による海水汚濁の判定。海洋科学，**2** (9/10)，33—39。
 [Kobayashi, Naomasa: Evaluation of marine pollution using fertilized sea urchin eggs. Marine Sciences Monthly, **2** (9/10), 33—39.]
- *森野浩：ベニボヤの体内より見出された端脚類 *Leucothoe spinicarpa* (Abildgaard) について (予報)。南紀生物，**12** (1)，8—10。
 [Morino, Hiroshi: Preliminary note on an amphipod *Leucothoe spinicarpa* (Abildgaard) found in the ascidian *Herdmania momus* (Savigny). Nanki Seibutu, **12** (1), 8—10.] (Jap. **187**)
- Nakayama, Yumiko: Successive changes of the vertical distribution of *Aidanosagitta delicata* (Tokioka) at a fixed station in a cove of Tanabe Bay. Publ. Seto Mar. Biol. Lab., **18** (3), 207—213. **526**
- Nishimura, Saburo [西村三郎]: Recent records of Baird's beaked whale in the Japan Sea. Ibid., **18** (1), 61—68. **522**
- *——：日本海の沿岸生物相をめぐるひとつの考え。南紀生物，**12** (2)，34—38。
 [Nishimura, Saburo: An idea concerning the littoral biota of the Japan Sea. Nanki Seibutu, **12** (2), 34—38.] (Jap. **188**)
- *——：海産動物の生育回遊における豊産への適応性。哺乳類科学，nos. 20/21 (徳田御稔先生退官記念号)，121—130。
 [——： Adaptations to a high fecundity as observed in the ontogenetic migration of marine animals. Mammal Sciences, nos. 20/21 (Commemoration volume on the retirement of Dr. Mitoshi Tokuda), 121—130.] (Jap. **189**)
- 岡田 稔・田原 胖：バフンウニにおける二次性徴の発生。動雑，**79** (2)，46—52。
 [Okada, Minoru & Tahara, Yutaka: Normal development of secondary sexual characters in the sea urchin, *Hemicentrotus pulcherrimus*. Zool. Mag., **79** (2), 46—52.] (Jap. **186**)
- Senawong, Chokechai: Preliminary observations on the moving ability of a littoral mussel, *Hormomya mutabilis* (Gould). Publ. Seto Mar. Biol. Lab., **18** (4), 233—242. **530**
- Tokioka, Takasi [時岡 隆]: Contributions to Japanese ascidian fauna XXV. Notes on the variations in *Botrylloides violaceus* Oka, with the description of a new subspecies *tenuicoecus*. Ibid., **18** (1), 57—59. **521**
- ：Ascidians from Mindoro Island, the Philippines. Ibid., **18** (2), 75—107. **523**
- ：Droplets from the plankton net XXV. Record of a *Rostraria* from Seto. Ibid., **18** (4), 275—278. **532**
- *——：沿岸における自然保護について。沿岸海洋研究ノート，**8** (2)，33—36。
 [——： Nature conservation in Japanese coastal waters, especially in view of sea fisheries. Bull. Coastal Oceanogr., **8** (2), 33—36.] (Jap. **190**)
- & Nishimura, Saburo: Record of an unusually large specimen of *Thysanostoma thysanura* Haeckel (Scyphozoa: Rhizostomae). Publ. Seto Mar. Biol. Lab., **17** (6), 429—437. **519**
- Utinomi, Huzio: Studies on the cirripedian fauna of Japan IX. Distributional survey of thoracic cirripeds in the southeastern part of the Japan Sea. Ibid., **17** (5), 339—372, pl. XVIII. **517**

| | |
|--|------------|
| ———: New and rare commensal pedunculate cirripeds from Amakusa islands, western Kyushu, Japan. Ibid., 18 (3), 157–167. | 524 |
| Werner, Bernhardt: Contribution to the evolution in the genus <i>Stephanoscyphus</i> (Scyphozoa Coronatae) and ecology and regeneration qualities of <i>Stephano-</i> <i>scyphus racemosus</i> Komai. Ibid., 18 (1), 1–20 | 520 |
| Yoshida, Keisei: Studies on germling development and life-history in Ulvaceae and Monostromaceae. Part I. Ibid., 17 (6), 403–428. | 518 |
| ———: Ditto. Part II. Ibid., 18 (4), 243–273. | 531 |

* Japanese text only.